



EBI Generic DB Entry Retrieval

```

ID  CBBONT1      standard; DNA; PRO; 4199 BP.
XX
AC  L35496;
XX
SV  L35496.1
XX
DT  23-AUG-1994 (Rel. 40, Created)
DT  04-MAR-2000 (Rel. 63, Last updated, Version 3)
XX
DE  Clostridium botulinum proteolytic type F neurotoxin bont/F gene, complete
DE  cds, and nontoxic-nonhaemagglutinin (ntnH) gene, 3' end.
XX
KW  bont/F gene; neurotoxin type F; nontoxic-nonhaemagglutinin; ntnH gene.
XX
OS  Clostridium botulinum
OC  Bacteria; Firmicutes; Clostridia; Clostridiales; Clostridiaceae;
OC  Clostridium.
XX
RN  [1]
RP  1-4199
RA  Elmore M.J., Bodsworth N.J., Whelan S.M., Minton N.P.;
RT  "The complete nucleotide sequence of the gene coding for proteolytic type
RT  neurotoxin of Clostridium botulinum";
RL  Unpublished.
XX
DR  SPTREMBL; Q45850; Q45850.
DR  SPTREMBL; Q57236; Q57236.
XX
FH  Key          Location/Qualifiers
FH
FT  source      1..4199
FT                /db_xref="taxon:1491"
FT                /organism="Clostridium botulinum"
FT  CDS        <1..171
FT                /codon_start=1
FT                /db_xref="SPTREMBL:Q45850"
FT                /note="putative"
FT                /transl_table=11
FT                /gene="ntnH"
FT                /product="nontoxic-nonhaemagglutinin"
FT                /protein_id="AAA23209.1"
FT                /translation="KKMIISNNIFISNCLTLSCGGKYICLSMKDENYNWMICNNESNI
FT                KKAYLWTLKEV"
FT  CDS        182..4018
FT                /codon_start=1
FT                /db_xref="SPTREMBL:Q57236"
FT                /note="putative"

```

```

FT      /transl_table=11
FT      /gene="bonT/F"
FT      /function="vertebrate neurotoxin"
FT      /product="neurotoxin type F"
FT      /protein_id="AAA23210.1"
FT      /translation="MPVVINSFNNDPVNDDTILYMQIPYEEKSKKYYKAFEIMRNVW
FT      IPERNTIGTDPSPDFPPASLENGSSAYYDPNYLTDAEKDRLKTTIKLFKRINSNPA
FT      EVLLQEISYAKPYLGNEHTPINEFHPVTRTTSVNIKSSTNVKSSIIINLLVLGAGPDI
FT      ENSSYPVRKLMDSGGVYDPSNDGFGSINIVTFSPEYEYTFNDISGGYNSSTESFIADP
FT      ISLAHELIHALHGLYGARGVTYKETIKVKQAPLMIAEKPIRLEEFLTFGGQDLNIITS
FT      MKEKIYNNLLANYEKIATRLSRVNSAPPEYDINEYKDYFQWKYGLDKNADGSYTVNEN
FT      FNEIYKKLYSFTEIDLANKFKVKCRNTYFIKYGFLKVPNLLDDDIYTVSEGFNIGNLA
FT      NNRGQNIKLNPKIIDSIPDKGLVEKIVKFKSVIPRKGTAPPRLCIRVNNRELFVA
FT      ESSYNENDINTPKEIDDTNLNNYRNNDDEVILDYNSETIPQISNQTLNLTQVQDDSY
FT      PRYDSNGTSEIEHNVDLNVFFYLHAQKVPEGETNISLTSSIDTALSEESQVYTFFS
FT      EFINTINKPVHAALFISWINQVIRDFTEATQKSTFDKIADISLVVPYVGLALNIGNE
FT      QKENFKEAFELLAGILLEFVPELLIPTILVFTIKSFIGSSENKNIKAINNSIMER
FT      TKWKEIYSWIVSNWLTRINTQFNKRKEQMYQALQNVDAIKTVIEYKNNYTSDERNR
FT      ESEYNINNIREELNKKVSLAMENIERFITESSIFYLMKLINEAKVSKLREYDEGVKEY
FT      LDYISEHRSILGNSVQELNDLVTSTLNNSIPFELSSYTNDKILILYFNKLYKKIKDNS
FT      LDMRYENNKFIDISGYGSNISINGDVYIYSTNRNQFGIYSSKPEVNIAQNNDIIYNG
FT      YQNFESISFWVRIPKYFNKVNLNNEYTIIDCIRNNNSGWKISLNYNKIITWLQDTAGNN
FT      KLVFNYSQTMISISDYINKWIFVTITNNRLGNSRIYINGNLIDEKISINLGDIVSDNI
FT      FKIVGCNDTRYVGIRYFKVFDTELGKTEIETLYSDEPDPSILKDFWGNLYLLNKRYYL
FT      NLLRTDKSITQNSNFLNINQQRGVYQKPNIFSNTRLYTGVEVIRKNGSTDISNTDNF
FT      RKNDLAYINVVDREVEYRLYADISIAKPEKIKLIRTSNSNSNLGQIIVMDSIGNNCT
FT      NFQNNNGGNIGLLGFHSNNLVASSWYNNIRKNTSSNGCFWSFISKEHGWQEN"
XX
SQ

```

Sequence 4199 BP; 1731 A; 432 C; 659 G; 1377 T; 0 other;

aagaagatga	taattagtaa	taatatattt	atttccaatt	gtttaactct	atcttgtggc	6
ggtaaatata	tatgtttatc	tatgaaagat	gaaaactata	attggatgat	atgtaataat	12
gaaagcaaca	tacctaaaa	ggcatattta	tggacattga	aagaagtata	ggggggattt	18
tatgccagtt	gtaataaata	gttttaatta	taatgacctt	gttaatgatg	atacaatttt	24
atacatgcag	ataccatatg	aagaaaaaag	taaaaaatat	tataaagctt	ttgagattat	30
gcgtaattgt	tggataattc	ctgagagaaa	tacaatagga	acggatccta	gtgattttga	36
tccaccggct	tcattagaga	acggaagcag	tgcttattat	gatcctaatt	atttaaccac	42
tgatgctgaa	aaagatagat	atttaaaaac	aacgataaaa	ttatttaaga	gaattaatag	48
taatcctgca	ggggaagtgt	tgttacaaga	aatatcatat	gctaaacctt	atttaggaaa	54
tgaacacacg	ccaattaatg	aattccatcc	agttactaga	actacaagtg	ttaatatata	60
atcatcaact	aatgttaaaa	gttcaataat	attgaatctt	cttgatttgg	gagcaggacc	66
tgatatattt	gaaaattctt	cttaccctgt	tagaaaaacta	atggattcag	gtggagttta	72
tgacccaagt	aatgatggtt	ttggatcaat	taatatcggt	acattttcac	ctgaatatga	78
atatactttt	aatgatatta	gtggagggtta	taacagtagt	acagaatcat	ttattgcaga	84
tcctgcaatt	tcactagctc	atgaattgat	acatgcactg	catggattat	acggggctag	90
gggagttact	tataaagaga	ctataaaaag	aaagcaagca	cctcttatga	tagccgaaaa	96
acccataagg	ctagaagaat	ttttaacctt	tggaggtcag	gatttaaata	ttattactag	102
tgctatgaag	gaaaaaatat	atacaaatct	tttagctaac	tatgaaaaaa	tagctactag	108
acttagtaga	gttaaatagtg	ctcctcctga	atatgatatt	aatgaatata	aagattattt	114
tcaatggaag	tatgggctag	ataaaaaatgc	tgatggaagt	tatactgtaa	atgaaaaata	120
atttaatgaa	atttataaaa	aattatatag	ctttacagag	attgacttag	caaataaatt	126
taaagtaaaa	tgtagaaata	cttattttat	taaatatgga	tttttaaaag	ttccaaaattt	132
gtagatgat	gatattttata	ctgtatcaga	ggggtttaat	ataggtaatt	tagcagtaaa	138
caatcgcgga	caaaatataa	agttaaatcc	taaaattatt	gattccattc	cagataaagg	144
tctagtggaa	aagatcggtta	aattttgtta	gagcgttatt	cctagaaaag	gtacaaaagg	150
gccaccgcga	ctatgcatta	gagtaataaa	tagggagtta	ttttttgtag	cttcagaaaag	156
tagctataat	gaaaatgata	ttaatacacc	taaagaaatt	gacgatacaa	caaatctaaa	162
taataattat	agaaataaatt	tagatgaagt	tatttttagat	tataatagtg	agacaatacc	168
tcaaatatca	aatcaaacat	taaatacact	tgtacaagac	gatagttatg	tgccaagata	174
tgattctaag	ggaacaagtg	aaatagagga	acataatggt	gttgacctta	atgtattttt	180
ctattttacat	gcacaaaaag	taccagaagg	tgaactaat	ataagtttaa	cttcttcaat	186
tgatacggca	ttatcagaag	aatcgcaagt	atatacattc	ttttcttcag	agtttattaa	192
tactatcaat	aaacctgtac	acgcagcact	atttataagt	tggataaatc	aagtaataag	198
agattttact	actgaagcta	cacaaaaaag	tacttttgat	aagattgcag	acatatcttt	204
agttgtacca	tatgtaggtc	ttgcttttaa	tataggtaat	gaggtacaaa	aagaaaattt	210
taaggaggca	tttgaattat	taggagcggt	tattttatta	gaatttgtgc	cagagctttt	216

```
aattcctaca attttagtgt ttacaataaa atcctttata ggttcatctg agaataaaaa 222
taaaatcatt aaagcaataa ataattcatt aatggaaaga gaaacaaagt ggaagaaat 228
atatagttgg atagtatcaa attggcttac tagaattaat acacaattta ataaaagaaa 234
agaacaaatg tatcaagctt tgcaaaatca agtagatgca ataaaaacag taatagaata 240
taaatataat aattatactt cagatgagag aaatagactt gaatctgaat ataatatcaa 246
taatataaga gaagaattga acaaaaaagt ttcttttagca atggaaaata tagagagatt 252
tataacagag agttctatat tttatttaat gaagttaata aatgaagcca aagttagtaa 258
attaagagaa tatgatgaag gcgttaagga atatttgcta gactatattt cagaacatag 264
atcaatttta ggaaatagtg tacaagaatt aaatgattta gtgactagta ctctgaataa 270
tagtattcca ttggaacttt cttcatatac taatgataaa attctaattt tatattttta 276
taaattatat aaaaaaatta aagataactc tatttttagat atgcgatatg aaaataataa 282
atztatagat atctctggat atggttcaaa tataagcatt aatggagatg tatataattta 288
ttcaacaaat agaaatcaat ttggaatata tagtagtaag cctagtgaag ttaatatagc 294
tcaaaataat gatattatat acaatggtag atatcaaaat tttagtatta gtttctgggt 300
aaggattcct aaatacttca ataaagtga tcttaataat gaatatacta taatagattg 306
tataaggaat aataattcag gatggaaaat atcacttaat tataataaaa taatttggac 312
tttacaagat actgctggaa ataatacaaaa actagttttt aattatacac aaatgattag 318
tatatctgat tatataaata aatggatttt tgtactattt actaataata gattaggcaa 324
ttctagaatt tacatcaatg gaaatttaat agatgaaaaa tcaatttcga atttaggtga 330
tattcatgtt agtgataata tattatttaa aattgttggt tgtaatgata caagatatgt 336
tggtataaga tattttaaag tttttgatac ggaattaggt aaaacagaaa ttgagacttt 342
atatagtgat gagccagatc caagtatctt aaaagacttt tggggaaatt atttggtata 348
taataaaaaga tattatttat tgaatttact aagaacagat aagtctatta ctcaaatc 354
aaactttcta aatattaatc aacaaagagg tgtttatcag aaaccaaata ttttttccaa 360
cactagatta tatacaggag tagaagtta tataagaaaa aatggatcta cagatatatc 366
taatacagat aattttgtta gaaaaaatga tctggcatat attaatgtag tagatcgtga 372
tgtagaatat cggctatatg ctgatatatc aattgcaaaa ccagagaaaa taataaaatt 378
aataagaaca tctaattcaa acaatagctt aggtcaaatt atagttatgg attcaatagg 384
aaataattgc acaatgaatt ttcaaaacaa taatgggggc aatataggat tactaggttt 390
tcattcaaat aatttggttg ctagtagttg gtattataac aatatacgaa aaaatactag 396
cagtaatgga tgcttttgga gttttatttc taaagagcat ggatggcaag aaaactaata 402
taataattca aaaaataggt attaaaatag aggtaatata tattaccctc tattttggaa 408
taattttaat atattatatg aaacatatat aaatttaaag ataatatata atcaagacac 414
aaattcaaat tagaaatata aaatgaagta aatgaaaagt gtaaaaagtc attaaataa 419
```

//

Please contact support@ebi.ac.uk with any problems or suggestions regarding this site.

L1 ANSWER 13 OF 17 MEDLINE
 AN 97016817 MEDLINE
 DN 97016817 PubMed ID: 8863443
 TI Organization and phylogenetic interrelationships of genes encoding components of the botulinum toxin complex in proteolytic *Clostridium botulinum* types A, B, and F: evidence of chimeric sequences in the gene encoding the nontoxic nonhemagglutinin component.
 AU East A K; Bhandari M; Stacey J M; Campbell K D; Collins M D
 CS Department of Microbiology, Institute of Food Research, Reading, Berkshire, United Kingdom.. alison.east@bbsrc.ac.uk
 SO INTERNATIONAL JOURNAL OF SYSTEMATIC BACTERIOLOGY, (1996 Oct) 46 (4) 1105-12.
 Journal code: 0042143. ISSN: 0020-7713.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 OS GENBANK-X78230; GENBANK-X87974; GENBANK-X92973; GENBANK-X96491; GENBANK-X96492; GENBANK-X96493; GENBANK-X96494
 EM 199611
 ED Entered STN: 19961219
 Last Updated on STN: 19980206
 Entered Medline: 19961115
 AB The cluster of genes encoding components of the **botulinum neurotoxin** (BoNT) complex was mapped in proteolytic (group I) *Clostridium botulinum* strains encoding BoNT types A, B, and F. Two different arrangements of genes were found: type A strain 62A and type B strain NCTC 7273 have similar organizations of genes encoding BoNT, the nontoxic nonhemagglutinin component (NTNH), hemagglutinin components, and P-21; **type F** strain Langeland has genes encoding BoNT, NTNH, and P-21, and a previously unidentified open reading frame encoding a protein of 416 amino acids. A group of type A strains typified by infant strain Kyoto-F, which is unlike type A strain 62A, lacks genes for hemagglutinin components and exhibits an organization similar to that of **type F**. Sequencing and pairwise analysis revealed the presence of possible chimeric sequences in some NTNH genes of proteolytic *C. botulinum*. Discordance in genealogical trees derived from different regions of the NTNH genes was observed which could be symptomatic of recombination and which may indicate that the NTNH gene represents a hot spot for such events within the cluster of genes encoding the BoNT complex. It is also evident that the phylogenetics of the NTNH gene, which is linked to the gene encoding BoNT, does not mirror the evolutionary history of the BoNT, upon which the *C. botulinum* species complex is defined and subdivided.

L1 ANSWER 14 OF 17 MEDLINE
 AN 94297488 MEDLINE
 DN 94297488 PubMed ID: 7764998
 TI Conserved structure of genes encoding components of **botulinum neurotoxin** complex M and the sequence of the gene coding for the nontoxic component in nonproteolytic *Clostridium botulinum* **type F**.
 AU East A K; Collins M D
 CS Department of Microbiology, Institute of Food Research, Reading Laboratory, UK.
 SO CURRENT MICROBIOLOGY, (1994 Aug) 29 (2) 69-77.
 Journal code: 7808448. ISSN: 0343-8651.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Biotechnology
 EM 199408

ED Entered STN: 19950809
 Last Updated on STN: 19970203
 Entered Medline: 19940810

AB For investigation of the genes of proteins associated in vivo with **botulinum neurotoxin** (BoNT), polymerase chain reaction (PCR) experiments were carried out with oligonucleotide primers designed to regions of the nontoxic-nonhemagglutinin (NTNH) gene of *Clostridium botulinum* type C. The primers were used to amplify a DNA fragment from genomic DNA of *C. botulinum* types A, B, E, F, G and toxigenic strains of *Clostridium barati* and *Clostridium butyricum*. The amplified product from all of these strains hybridized with an internal oligonucleotide probe, whereas all nontoxigenic clostridia tested gave no PCR product and showed no reaction with the probe. The NTNH gene was shown to be located upstream of the gene encoding BoNT, thereby revealing a conserved structure for genes encoding the proteins of the M complex of the progenitor botulinum toxin in these organisms. The sequence of the NTNH gene of nonproteolytic *C. botulinum* **type F** was determined by PCR amplification and sequencing of overlapping cloned fragments. NTNH/F showed 71% and 61% identity with NTNH of *C. botulinum* type E and type C respectively.

L1 ANSWER 15 OF 17 MEDLINE
 AN 93012902 MEDLINE
 DN 93012902 PubMed ID: 1398040
 TI Sequence of the gene encoding **type F** neurotoxin of *Clostridium botulinum*.
 AU East A K; Richardson P T; Allaway D; Collins M D; Roberts T A; Thompson D E
 CS Department of Microbiology, AFRC Institute of Food Research, Reading, UK.
 SO FEMS MICROBIOLOGY LETTERS, (1992 Sep 15) 75 (2-3) 225-30.
 Journal code: 7705721. ISSN: 0378-1097.
 CY Netherlands
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 OS GENBANK-M92906
 EM 199211
 ED Entered STN: 19930122
 Last Updated on STN: 19970203
 Entered Medline: 19921119

AB Primers designed to conserved regions of botulinum and tetanus clostridial toxins were used to amplify DNA fragments from non-proteolytic *Clostridium botulinum* **type F** (202F) DNA using polymerase chain reaction technology. The fragments were cloned and the complete nucleotide sequence of the gene encoding **type F** toxin determined. Analysis of the nucleotide sequence demonstrated the presence of an open frame encoding a protein of 1274 amino acids, similar to other **botulinum neurotoxins**. Upstream of the toxin gene is the end of an open reading frame which encodes the C-terminus of a protein with homology to non-toxic-non-hemagglutinin component of type C progenitor toxin.

L1 ANSWER 16 OF 17 MEDLINE
 AN 90262533 MEDLINE
 DN 90262533 PubMed ID: 2188647
 TI Botulinum **type F** neurotoxin. Large-scale purification and characterization of its binding to rat cerebrocortical synaptosomes.
 AU Wadsworth J D; Desai M; Tranter H S; King H J; Hambleton P; Melling J; Dolly J O; Shone C C
 CS Department of Biochemistry, Imperial College of Science and Technology, London, U.K.
 SO BIOCHEMICAL JOURNAL, (1990 May 15) 268 (1) 123-8.

Journal code: 2984726R. ISSN: 0264-6021.

CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199006
 ED Entered STN: 19900720
 Last Updated on STN: 19970203
 Entered Medline: 19900628

AB 1. A large-scale purification procedure has been developed for Clostridium botulinum **type F** neurotoxin. Commencing with 160 litres of bacterial culture, 101 mg of purified **type F** neurotoxin with a specific toxicity of 2×10^7 mouse LD50 (median lethal dose).mg-1 were obtained. 2. Purified **type F** neurotoxin was labelled to high specific radioactivity (900-1360 Ci/mmol) without loss of biological activity using a chloramine-T procedure. Of the two neurotoxin subunits, the heavy chain was preferentially radiolabelled. 3. Radiolabelled **type F** neurotoxin displayed specific saturable binding to rat synaptosomes. At least two pools of acceptors were evident: a low content of high-affinity acceptors sites [KD approximately 0.15 nM; Bmax (maximal binding) 20 fmol/mg] and a larger pool of lower-affinity sites (KD greater than 20 nM; Bmax greater than 700 fmol/mg). Both pools of acceptors were sensitive to trypsin and neuraminidase treatment, which suggests that protein and sialic acid residues are components of the synaptosomal acceptors. 4. Experiments investigating competition among **botulinum neurotoxin** types A, B, E and F for acceptors on rat brain synaptosomes showed that **type F** neurotoxin binds to acceptor molecules which are completely distinct from those of the other three neurotoxins.

L1 ANSWER 17 OF 17 MEDLINE
 AN 84018414 MEDLINE
 DN 84018414 PubMed ID: 6353671
 TI Amino acid composition of Clostridium botulinum **type F** neurotoxin.
 AU DasGupta B E; Rasmussen S
 NC NS 17742 (NINDS)
 SO TOXICON, (1983) 21 (4) 566-9.
 Journal code: 1307333. ISSN: 0041-0101.
 CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 198311
 ED Entered STN: 19900319
 Last Updated on STN: 19970203
 Entered Medline: 19831123

AB To develop reliable data on the amino acid composition of **type F botulinum neurotoxin**, three batches of the neurotoxin were analyzed. Each batch was isolated from a separate neurotoxin producing bacterial culture. Two batches had inoculum from one source and the other batch one from a different source. Two batches of the neurotoxin were purified by the same method and one was purified by a different method. The neurotoxin preparations were found comparable in purity and similar in amino acid composition. The best estimate of number of amino acid residues per neurotoxin molecule (mol. wt. 155,000) was: Asp218 Thr80 Ser105 Glu128 Pro47 Gly69 Ala47 Val72 Cys9 Met14 Ile128 Leu104 Tyr86 Phe60 Lys90 His13 Arg51 Trp23.

=>

L2 ANSWER 7 OF 10 MEDLINE
 AN 95179082 MEDLINE
 DN 95179082 PubMed ID: 7874084
 TI Inhibition of neurotransmitter release by clostridial neurotoxins correlates with specific proteolysis of synaptosomal proteins.
 AU Blasi J; Binz T; Yamasaki S; Link E; Niemann H; Jahn R
 CS Departament de Biologia Cel.lular i Anatomia Patologica, Facultat de Medicina, Universitat de Barcelona, Spain.
 SO JOURNAL OF PHYSIOLOGY, PARIS, (1994) 88 (4) 235-41.
 Journal code: 9309351. ISSN: 0928-4257.
 CY France
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199504
 ED Entered STN: 19950419
 Last Updated on STN: 19970203
 Entered Medline: 19950404
 AB Rat brain synaptosomes were used to study the effect of several clostridial neurotoxins on the neurotransmitter release. In this system the blockade of transmitter release correlated with the proteolytic activity of the toxins. Blockade of glutamate release was linked to selective proteolysis of one of the following synaptic proteins: synaptobrevin (BoNT/D, **BoNT/F**); SNAP-25 (BoNT/A, BoNT/E), or HPC-1/syntaxin (BoNT/C1). All the toxins used had an inhibitory effect on synaptosomes with the exception of **BoNT/F**. **BoNT/F** cleaved synaptobrevin in permeabilized synaptosomes but failed to produce the same effect on intact synaptosomes.

L2 ANSWER 8 OF 10 MEDLINE
 AN 94230352 MEDLINE
 DN 94230352 PubMed ID: 8175689
 TI Cleavage of members of the synaptobrevin/VAMP family by types D and F botulinal neurotoxins and tetanus toxin.
 AU Yamasaki S; Baumeister A; Binz T; Blasi J; Link E; Cornille F; Roques B; Fykse E M; Sudhof T C; Jahn R; +
 CS Department of Microbiology, Federal Research Center for Virus Diseases of Animals, Tübingen, Federal Republic of Germany.
 SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1994 Apr 29) 269 (17) 12764-72.
 Journal code: 2985121R. ISSN: 0021-9258.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199406
 ED Entered STN: 19940620
 Last Updated on STN: 19970203
 Entered Medline: 19940606
 AB Tetanus toxin (TeTx) and the various forms of botulinal neurotoxins (BoNT/A to BoNT/G) potently inhibit neurotransmission by means of their L chains which selectively proteolyze synaptic proteins such as synaptobrevin (TeTx, BoNT/B, **BoNT/F**), SNAP-25 (BoNT/A), and syntaxin (BoNT/C1). Here we show that BoNT/D cleaves rat synaptobrevin 1 and 2 in toxified synaptosomes and in isolated vesicles. In contrast, synaptobrevin 1, as generated by in vitro translation, is only a poor substrate for BoNT/D, whereas this species is cleaved by **BoNT/F** with similar potency. Cleavage by BoNT/D occurs at the peptide bond Lys59-Leu60 which is adjacent to the **BoNT/F** cleavage site (Gln58-Lys59) and again differs from the site hydrolyzed by TeTx and BoNT/B (Gln76-Phe77). Cellubrevin, a recently

discovered isoform expressed outside the nervous system, is efficiently cleaved by all three toxins examined. For further characterization of the substrate requirements of BoNT/D, we tested amino- and carboxyl-terminal deletion mutants of synaptobrevin 2 as well as synthetic peptides. Shorter peptides containing up to 15 amino acids on either side of the cleavage site were not cleaved, and a peptide extending from Arg47 to Thr116 was a poor substrate for all three toxins tested. However, cleavability was restored when the peptide is further extended at the NH2 terminus (Thr27-Thr116) demonstrating that NH2 terminally located sequences of synaptobrevin which are distal from the respective cleavage sites are required for proteolysis. To further examine the isoform specificity, several mutants of rat synaptobrevin 2 were generated in which individual amino acids were replaced with those found in rat synaptobrevin 1. We show that a Met46 to Ile46 substitution drastically diminishes cleavability by BoNT/D and that the presence of Val76 instead of Gln76 dictates the reduced cleavability of synaptobrevin isoforms by TeTx.

L2 ANSWER 9 OF 10 MEDLINE
 AN 94013372 MEDLINE
 DN 94013372 PubMed ID: 8408542
 TI Gene probes for identification of the botulinum neurotoxin gene and specific identification of neurotoxin types B, E, and F.
 AU Campbell K D; Collins M D; East A K
 CS Reading Laboratory, Institute of Food Research, Agriculture and Food Research Council, United Kingdom.
 SO JOURNAL OF CLINICAL MICROBIOLOGY, (1993 Sep) 31 (9) 2255-62.
 Journal code: 7505564. ISSN: 0095-1137.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 OS GENBANK-X70814; GENBANK-X70815; GENBANK-X70816; GENBANK-X70817; GENBANK-X70818; GENBANK-X70819; GENBANK-X70820; GENBANK-X70821
 EM 199311
 ED Entered STN: 19940117
 Last Updated on STN: 19970203
 Entered Medline: 19931102
 AB A polymerase chain reaction method was developed for the specific detection of the botulinum neurotoxin (BoNT) gene of Clostridium botulinum. Degenerate oligonucleotide primers, designed from the nucleotide sequence of the heavy chain of the BoNT gene, amplified a specific fragment of approximately 1.1 kb from strains of C. botulinum toxin types A, B, E, F, and G and neurotoxin-producing strains of Clostridium barati and Clostridium butyricum, but no fragment was obtained from nontoxicogenic strains. The fragments amplified from several strains of C. botulinum types B, E, and F were cloned in Escherichia coli and their nucleotide sequences were determined. Sequences within this region were used to design oligonucleotide probes specific for BoNT type B (BoNT/B), BoNT/E, and **BoNT/F** genes. An additional probe was designed for the detection of the **BoNT/F** gene of C. barati, which differed in sequence from **BoNT/F** genes of both proteolytic and nonproteolytic strains of C. botulinum.

L2 ANSWER 10 OF 10 MEDLINE
 AN 93012902 MEDLINE
 DN 93012902 PubMed ID: 1398040
 TI Sequence of the gene encoding type F neurotoxin of Clostridium botulinum.
 AU East A K; Richardson P T; Allaway D; Collins M D; Roberts T A; Thompson D E
 CS Department of Microbiology, AFRC Institute of Food Research, Reading, UK.
 SO FEMS MICROBIOLOGY LETTERS, (1992 Sep 15) 75 (2-3) 225-30.
 Journal code: 7705721. ISSN: 0378-1097.

CY Netherlands
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
OS GENBANK-M92906
EM 199211
ED Entered STN: 19930122
Last Updated on STN: 19970203
Entered Medline: 19921119
AB Primers designed to conserved regions of botulinum and tetanus clostridial toxins were used to amplify DNA fragments from non-proteolytic Clostridium botulinum type F (202F) DNA using polymerase chain reaction technology. The fragments were cloned and the complete nucleotide sequence of the gene encoding type F toxin determined. Analysis of the nucleotide sequence demonstrated the presence of an open frame encoding a protein of 1274 amino acids, similar to other botulinum neurotoxins. Upstream of the toxin gene is the end of an open reading frame which encodes the C-terminus of a protein with homology to non-toxic-non-hemagglutinin component of type C progenitor toxin.

=>



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Boc
Search		Protein	for				Go	Clear
		Limits	Preview/Index	History	Clipboard		Details	
Display	default	Save	Text	Add to Clipboard	Get Subsequence			

☐ 1: P10845. Botulinum neuroto...[gi:399133]

[Links](#)

LOCUS BXAI_CLOBO 1296 aa linear BCT 15-JUN-2002
 DEFINITION Botulinum neurotoxin type A precursor (BoNT/A) (Bontoxilysin A) (BOTOX) [Contains: Botulinum neurotoxin A, light-chain; Botulinum neurotoxin A, heavy-chain].
 ACCESSION P10845
 VERSION P10845 GI:399133
 DBSOURCE swissprot: locus BXAI_CLOBO, accession P10845; class: standard. extra accessions: P18639, P01561, created: Jul 1, 1989. sequence updated: Jul 1, 1993. annotation updated: Jun 15, 2002. xrefs: gi: 40381, gi: 40382, gi: 144864, gi: 144865, gi: 1619251, gi: 1619253, gi: 2160224, gi: 2160225, gi: 144880, gi: 551776, gi: 279630, gi: 98562, pdb accession 3BTA xrefs (non-sequence databases): MEROPS M27.002, InterPro IPR000395, InterPro IPR000130, Pfam PF01742, PRINTS PR00760, ProDom PD001963, PROSITE PS00142
 KEYWORDS Neurotoxin; Transmembrane; Hydrolase; Metalloprotease; Zinc; Pharmaceutical; 3D-structure.
 SOURCE Clostridium botulinum.
 ORGANISM Clostridium botulinum Bacteria; Firmicutes; Clostridia; Clostridiales; Clostridiaceae; Clostridium.
 REFERENCE 1 (residues 1 to 1296)
 AUTHORS Thompson, D.E., Brehm, J.K., Oultram, J.D., Swinfield, T.J., Shone, C.C., Atkinson, T., Melling, J. and Minton, N.P.
 TITLE The complete amino acid sequence of the Clostridium botulinum type A neurotoxin, deduced by nucleotide sequence analysis of the encoding gene
 JOURNAL Eur. J. Biochem. 189 (1), 73-81 (1990)
 MEDLINE 90235864
 PUBMED 2185020
 REMARK SEQUENCE FROM N.A. STRAIN=NCTC 2916
 REFERENCE 2 (residues 1 to 1296)
 AUTHORS Binz, T., Kurazono, H., Wille, M., Frevert, J., Wernars, K. and Niemann, H.
 TITLE The complete sequence of botulinum neurotoxin type A and comparison with other clostridial neurotoxins
 JOURNAL J. Biol. Chem. 265 (16), 9153-9158 (1990)
 MEDLINE 90264400
 PUBMED 2160960
 REMARK SEQUENCE FROM N.A. STRAIN=62A
 REFERENCE 3 (residues 1 to 1296)
 AUTHORS East, A.K., Bhandari, M., Stacey, J.M., Campbell, K.D. and Collins, M.D.
 TITLE Organization and phylogenetic interrelationships of genes encoding components of the botulinum toxin complex in proteolytic Clostridium botulinum types A, B, and F: evidence of chimeric sequences in the gene encoding the nontoxic nonhemagglutinin

component

JOURNAL Int. J. Syst. Bacteriol. 46 (4), 1105-1112 (1996)

MEDLINE 97016817

PUBMED 8863443

REMARK SEQUENCE OF 1-65 FROM N.A.
STRAIN=62A

REFERENCE 4 (residues 1 to 1296)

AUTHORS Betley,M.J., Somers,E. and DasGupta,B.R.

TITLE Characterization of botulinum type A neurotoxin gene: delineation of the N-terminal encoding region

JOURNAL Biochem. Biophys. Res. Commun. 162 (3), 1388-1395 (1989)

MEDLINE 89350959

PUBMED 2669749

REMARK SEQUENCE OF 1-34 FROM N.A.
STRAIN=Hall

REFERENCE 5 (residues 1 to 1296)

AUTHORS Fujita,R., Fujinaga,Y., Inoue,K., Nakajima,H., Kumon,H. and Oguma,K.

TITLE Molecular characterization of two forms of nontoxic-nonhemagglutinin components of Clostridium botulinum type A progenitor toxins

JOURNAL FEBS Lett. 376 (1-2), 41-44 (1995)

MEDLINE 96096783

PUBMED 8521962

REMARK SEQUENCE OF 1-18 FROM N.A.
STRAIN=Type A NIH

REFERENCE 6 (residues 1 to 1296)

AUTHORS Schmidt,J.J., Sathyamoorthy,V. and DasGupta,B.R.

TITLE Partial amino acid sequence of the heavy and light chains of botulinum neurotoxin type A

JOURNAL Biochem. Biophys. Res. Commun. 119 (3), 900-904 (1984)

MEDLINE 84178501

PUBMED 6370252

REMARK SEQUENCE OF 1-16.

REFERENCE 7 (residues 1 to 1296)

AUTHORS Dasgupta,B.R., Foley,J. and Niece,R.

TITLE Partial sequence of the light chain of botulinum neurotoxin type A

JOURNAL Biochemistry 26, 4162-4162 (1987)

REMARK SEQUENCE OF 1-46.

REFERENCE 8 (residues 1 to 1296)

AUTHORS DasGupta,B.R. and Dekleva,M.L.

TITLE Botulinum neurotoxin type A: sequence of amino acids at the N-terminus and around the nicking site

JOURNAL Biochimie 72 (9), 661-664 (1990)

MEDLINE 91120847

PUBMED 2126206

REMARK SEQUENCE OF 1-5 AND 444-456.

REFERENCE 9 (residues 1 to 1296)

AUTHORS Sathyamoorthy,V., Dasgupta,B.R., Foley,J. and Niece,R.L.

TITLE Botulinum neurotoxin type A: cleavage of the heavy chain into two halves and their partial sequences

JOURNAL Arch. Biochem. Biophys. 266 (1), 142-151 (1988)

MEDLINE 89024662

PUBMED 3178218

REMARK SEQUENCE OF 448-464 AND 872-895.

REFERENCE 10 (residues 1 to 1296)

AUTHORS Shone,C.C., Hambleton,P. and Melling,J.

TITLE Inactivation of Clostridium botulinum type A neurotoxin by trypsin and purification of two tryptic fragments. Proteolytic action near the COOH-terminus of the heavy subunit destroys toxin-binding activity

JOURNAL Eur. J. Biochem. 151 (1), 75-82 (1985)

MEDLINE 85285016

PUBMED 3896784

REMARK SEQUENCE OF 448-482.

REFERENCE 11 (residues 1 to 1296)
AUTHORS Schiavo,G., Santucci,A., Dasgupta,B.R., Mehta,P.P., Jontes,J., Benfenati,F., Wilson,M.C. and Montecucco,C.
TITLE Botulinum neurotoxins serotypes A and E cleave SNAP-25 at distinct COOH-terminal peptide bonds
JOURNAL FEBS Lett. 335 (1), 99-103 (1993)
MEDLINE 94063091
PUBMED 8243676
REMARK IDENTIFICATION OF SUBSTRATE.
REFERENCE 12 (residues 1 to 1296)
AUTHORS Binz,T., Blasi,J., Yamasaki,S., Baumeister,A., Link,E., Sudhof,T.C., Jahn,R. and Niemann,H.
TITLE Proteolysis of SNAP-25 by types E and A botulinum neurotoxins
JOURNAL J. Biol. Chem. 269 (3), 1617-1620 (1994)
MEDLINE 94124495
PUBMED 8294407
REMARK IDENTIFICATION OF SUBSTRATE.
REFERENCE 13 (residues 1 to 1296)
AUTHORS Rigoni,M., Caccin,P., Johnson,E.A., Montecucco,C. and Rossetto,O.
TITLE Site-directed mutagenesis identifies active-site residues of the light chain of botulinum neurotoxin type A
JOURNAL Biochem. Biophys. Res. Commun. 288 (5), 1231-1237 (2001)
MEDLINE 21556941
PUBMED 11700044
REMARK MUTAGENESIS OF GLU-261; PHE-265 AND TYR-365.
REFERENCE 14 (residues 1 to 1296)
AUTHORS Lacy,D.B., Tepp,W., Cohen,A.C., DasGupta,B.R. and Stevens,R.C.
TITLE Crystal structure of botulinum neurotoxin type A and implications for toxicity
JOURNAL Nat. Struct. Biol. 5 (10), 898-902 (1998)
MEDLINE 98455071
PUBMED 9783750
REMARK X-RAY CRYSTALLOGRAPHY (3.3 ANGSTROMS).
COMMENT On or before Sep 14, 1993 this sequence version replaced gi:115193, gi:115174.

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. The original entry is available from <http://www.expasy.ch/sprot> and <http://www.ebi.ac.uk/sprot>

[FUNCTION] Inhibits acetylcholine release. The botulinum toxin binds with high affinity to peripheral neuronal presynaptic membrane, is then internalized by receptor-mediated endocytosis. The C-terminus of the heavy chain (H) is responsible for the adherence of the toxin to the cell surface while the N-terminus mediates transport of the light chain from the endocytic vesicle to the cytosol. After translocation, the light chain (L) hydrolyzes the 197-Gln-I-Arg-198 bond in SNAP-25, thereby blocking neurotransmitter release. Inhibition of acetylcholine release results in flaccid paralysis, with frequent heart or respiratory failure.

[CATALYTIC ACTIVITY] Limited hydrolysis of proteins of the neuroexocytosis apparatus, synaptobrevins, SNAP25 or syntaxin. No detected action on small molecule substrates.

[SUBUNIT] Disulfide-linked heterodimer of a light chain (L) and a heavy chain (H).

[SUBCELLULAR LOCATION] Secreted.

PHARMACEUTICAL: Available under the name BOTOX (Allergan) for the treatment of strabismus and blepharospasm associated with dystonia and cervical dystonia. Also used for the treatment of hemifacial spasm and a number of other neurological disorders characterized by abnormal muscle contraction.

[MISCELLANEOUS] There are seven antigenically distinct forms of

botulinum neurotoxin: Types A, B, C1, D, E, F, and G.
 [SIMILARITY] BELONGS TO PEPTIDASE FAMILY M27.
 [DATABASE] NAME=BOTOX product information Web site;
 WWW='http://www.botox.com/index.jsp?hp&productinfo'.
 [DATABASE] NAME=Protein Spotlight; NOTE=Issue 19 of February 2002;
 WWW='http://www.expasy.org/spotlight/articles/sptlt019.html'.

FEATURES	Location/Qualifiers
source	1..1296 /organism="Clostridium botulinum" /db_xref="taxon:1491"
gene	1..1296 /gene="BOTA" /note="BNA; ATX"
Protein	1..1296 /gene="BOTA" /product="Botulinum neurotoxin type A precursor" /EC_number="3.4.24.69"
Region	2..448 /gene="BOTA" /region_name="Mature chain" /note="BOTULINUM NEUROTOXIN A, LIGHT-CHAIN."
Region	2 /gene="BOTA" /region_name="Conflict" /note="P -> Q (IN REF. 1)."
Region	27 /gene="BOTA" /region_name="Variant" /note="V -> A."
Site	223 /gene="BOTA" /site_type="metal-binding" /note="ZINC (CATALYTIC)."
Site	224 /gene="BOTA" /site_type="active"
Site	227 /gene="BOTA" /site_type="metal-binding" /note="ZINC (CATALYTIC)."
Site	262 /gene="BOTA" /site_type="metal-binding" /note="ZINC (CATALYTIC)."
Site	262 /gene="BOTA" /site_type="mutagenized" /note="E->A: DRASTIC DECREASE IN ENZYMATIC ACTIVITY."
Site	266 /gene="BOTA" /site_type="mutagenized" /note="F->A: DECREASE IN ENZYMATIC ACTIVITY."
Site	366 /gene="BOTA" /site_type="mutagenized" /note="Y->A: DECREASE IN ENZYMATIC ACTIVITY."
Bond	bond(430,454) /gene="BOTA" /bond_type="disulfide" /note="INTERCHAIN."
Region	449..1296 /gene="BOTA" /region_name="Mature chain" /note="BOTULINUM NEUROTOXIN A, HEAVY-CHAIN."
Region	480

Revised: July 5, 2002.

ID Q57236 PRELIMINARY; PRT; 1278 AA.
 AC Q57236; Q45863;
 DT 01-NOV-1996 (TrEMBLrel. 01, Created)
 DT 01-NOV-1996 (TrEMBLrel. 01, Last sequence update)
 DT 01-OCT-2002 (TrEMBLrel. 22, Last annotation update)
 DE Botulinum neurotoxin type F (BONT/F protein).
 GN BONT/F.
 OS Clostridium botulinum.
 OC Bacteria; Firmicutes; Bacillus/Clostridium group; Clostridia;
 OC Clostridiales; Clostridiaceae; Clostridium.
 OX NCBI_TaxID=1491;
 RN [1]
 RP SEQUENCE FROM N.A.
 RC STRAIN=NCTC 10281;
 RA Hutson R.A., Collins M.D.;
 RL Submitted (AUG-1995) to the EMBL/GenBank/DDBJ databases.
 RN [2]
 RP SEQUENCE FROM N.A.
 RA Elmore M.J., Bodsworth N.J., Whelan S.M., Minton N.P.;
 RL Submitted (AUG-1994) to the EMBL/GenBank/DDBJ databases.
 RN [3]
 RP SEQUENCE OF 635-1000 FROM N.A.
 RC STRAIN=NCTC 1028;
 RX MEDLINE=94013372; PubMed=8408542;
 RA Campbell K., East A.K., Collins M.D.;
 RT "Gene probes for identification of the botulinal neurotoxin gene and
 RT specific identification of neurotoxin types B, E, and F."
 RL J. Clin. Microbiol. 31:2255-2262(1993).
 RN [4]
 RP SEQUENCE OF 1-27 FROM N.A.
 RC STRAIN=LANGELAND;
 RX MEDLINE=98404102; PubMed=9732534;
 RA East A.K., Bhandari M., Hielm S., Collins M.D.;
 RT "Analysis of the botulinum neurotoxin type F gene clusters in
 RT proteolytic and nonproteolytic Clostridium botulinum and Clostridium
 RT barati."
 RL Curr. Microbiol. 37:262-268(1998).
 DR EMBL; X81714; CAA57358.1; -.
 DR EMBL; L35496; AAA23210.1; -.
 DR EMBL; X70821; CAA50152.1; -.
 DR EMBL; X99064; CAA67512.1; -.
 DR HSSP; P10845; 3BTA.
 DR MEROPS; M27.002; -.
 DR InterPro; IPR000395; Bontoxilysin.
 DR InterPro; IPR006025; Zn_MTpeptdse.
 DR Pfam; PF01742; Peptidase_M27; 1.
 DR PRINTS; PR00760; BONTOXILYSIN.
 DR ProDom; PD001963; Bontoxilysin; 1.
 DR PROSITE; PS00142; ZINC_PROTEASE; UNKNOWN_1.
 KW Neurotoxin.
 SQ SEQUENCE 1278 AA; 147073 MW; A1BE1318431D6918 CRC64;
 MPVVINSFNY NDPVNDDTIL YMQIPYEEKS KKYYKAFEIM RNVWIIPERN TIGTDPSDFD
 PPASLENGSS AYYDPNYLTT DAEKDRYLKT TIKLFKRINS NPAGEVLLQE ISYAKPYLGN
 EHTPINEFHP VTRTTSVNIK SSTNVKSSII LNLLVLGAGP DIFENSSTYPV RKLMDSGGVY
 DPSNDGFGSI NIVTFSPEYE YTFNDISGGY NSSTESFIAD PAISLAHELI HALHGLYGAR
 GVTYKETIKV KQAPLMIAEK PIRLEEFLTF GGQDLNIITS AMKEKIYNNL LANYEKIATR
 LSRVNSAPE YDINEYKDYF QWKYGLDKNA DGSYTVNENK FNEIYKKLYS FTEIDLANKF
 KVKCRNTYFI KYGFLKVPNL LDDDIYTVSE GFNIGNLAVN NRGQNIKLNP KIIDSIPDKG
 LVEKIVKFECK SVIPRKGTKA PPRLCIRVNN RELFFVASES SYNENDINTP KEIDDTNIN
 NNYRNNLDEV ILDYNSETIP QISNQTLNLT VQDDSYVPRY DSNGTSEIEE HNVVDLNVFF
 YLHAQKVPEG ETNISLTSSI DTALSEESQV YTFFSSEFIN TINKPVHAAL FISWINQVIR
 DFTTEATQKS TFDKIADISL VVPYVGLALN IGNEVQKENF KEAFELLGAG ILLEFVPELL
 IPTILVFTIK SFIGSSENKN KIIKAINNSL MERETKWKEI YSWIVSNWLT RINTQFNKRK
 EQMYQALQNO VDAIKTVIEY KYNNTS DER NRLESEYNIN NIREELNKKV SLAMENIERF
 ITESSIFYLM KLINEAKVSK LREYDEGVKE YLLDYISEHR SILGNSVQEL NDLVTSTLNN

SIPFELSSYT NDKILILYFN KLYKKIKDNS ILDMRYENNK FIDISGYGSN ISINGDVYIY
STNRNQFGIY SSKPSEVNIA QNNDIIYNGR YQNFISFWV RIPKYFNKVN LNNEYTIIDC
IRNNNSGWKI SLNYNKIIWT LQDTAGNNQK LVFNYSQMS ISDYINKWIF VTITNNRLGN
SRIYINGNLI DEKSISNLGD IHVSDNLFK IVGCNDTRYV GIRYFKVFDI ELGKTEIETL
YSDEPDPSIL KDFWGNLYLY NKRYLLNLL RTDKSITQNS NFLNINQQRG VYQKPNIFSN
TRLYTGVEVI IRKNGSTDIS NTDNFVRKND LAYINVVDRI VEYRLYADIS IAKPEKIKL
IRTSNSNNSL GQIIVMDSIG NNCTMNFQNN NGGNIGLLGF HSNNLVASSW YYNNIRKNTS
SNGCFWSFIS KEHGWQEN

//